

The European Large Area *ISO* Survey: ELAIS

Seb Oliver¹, M. Rowan-Robinson, C. Cesarsky, L. Danese, A. Franceschini, R. Genzel, A. Lawrence, D. Lemke, R. McMahon, G. Miley, J-L. Puget, B. Rocca-Volmerange, P. Ciliegi, A. Efsthathiou, C. Gruppioni, P. Héraudeau, S. Serjeant, C. Surace and ELAIS consortium

¹*Imperial College of Science Technology and Medicine, Astrophysics Group, Blackett Laboratory, Prince Consort Rd., London, SW2 1BZ, s.oliver@ic.ac.uk*

²*To appear in 'Wide-field surveys in cosmology', Proc. XIV IAP meeting*

Abstract.

The European Large Area *ISO* Survey (ELAIS) has surveyed ~ 12 square degrees of the sky at $15\mu\text{m}$ and $90\mu\text{m}$ and subsets of this area at $6.75\mu\text{m}$ and $175\mu\text{m}$ using the Infrared Space Observatory (*ISO*). This project was the largest single open time programme executed by *ISO*, taking 375 hours of data. A preliminary catalogue of more than 1000 galaxies has been produced. In this talk we describe the goals of the project and present a provisional number count analysis at $15\mu\text{m}$.

1 Introduction

The European Large Area *ISO* Survey (ELAIS) is a collaboration involving 19 European institutes (in addition to the authors and others at their institutes the following people and others at their institutes are involved I. Gonzalez-Serrano, E. Kontizas, K. Mandolesi, J. Masegosa, K. Mattila, H. Norgaard-Nielsen, I. Perez-Fournon, M. Ward) and is the largest open time project being undertaken by *ISO*.

2 Key Scientific Goals

In this section we highlight some of the principal scientific motivations for this programme, though naturally there are many other goals which we have not space discuss.

The main extra-galactic population detected by IRAS was galaxies with high rates of star formation. These objects are now known to evolve with a strength comparable to AGN. The sensitivity of *ISO* allow us to detect these objects at much higher redshifts and thus obtain greater understanding of the cosmological evolution of star formation. This is directly complimentary to studies of star-formation history in the optical UV. Comparison of the star-formation rates determined in the FIR with that determined from the UV will give a direct estimate of the importance of dust obscuration, vitally important for models of cosmic evolution.

If elliptical galaxies underwent a massive burst of star-formation between $2 < z < 5$, they would be observable in the far infrared and may look like F10214. This survey will provide a powerful discrimination between such a top-down scenario and any hierarchical bottom-up merging model whose components are individually too faint to detect.

Table 1: Summary of ELAIS Survey Areas. These areas were selected primarily for having low Cirrus contamination, specifically $I_{100} < 1.5 \text{ MJy/sr}$ from the IRAS maps of Rowan-Robinson *et al.* (1991). For N1-3, S1 and all X areas we also restricted ourselves to regions of high visibility $> 25\%$ over the mission lifetime. For low Zodiacal background we required $|\beta| > 40$ and to avoid saturation of the CAM detectors we had to avoid any bright IRAS $12\mu\text{m}$ sources. The 6 additional small ($0.3^\circ \times 0.3^\circ$) rasters (X1-6) were centred on well studied areas of the sky or high- z objects

Name	Nominal Coordinates		M	N	ROLL	$\langle I_{100} \rangle$
	J2000		/ $^\circ$	/ $^\circ$	/ $^\circ$	/MJysr $^{-1}$
N1	$16^h 10^m 01^s$	$+54^\circ 30' 36''$	2.0	1.3	20	1.2
N2	$16^h 36^m 58^s$	$+41^\circ 15' 43''$	2.0	1.3	30	1.1
N3	$14^h 29^m 06^s$	$+33^\circ 06' 00''$	2.0	1.3	330	0.9
S1	$00^h 34^m 44^s$	$-43^\circ 28' 12''$	2.0	2.0	20	1.1
S2	$05^h 02^m 24^s$	$-30^\circ 35' 55''$	0.3	0.3	290	1.1

Unified models of AGN suggest that the central engine is surrounded by a dusty torus. The far infrared emission from the torus is much less sensitive to the viewing angle than the optical. Thus a far infrared selected sample of AGN will place important constraints on unification schemes.

IRAS uncovered a population with enormous far infrared luminosities, $L_{\text{FIR}} > 10^{12} L_\odot$. While most of these objects appear to have an AGN it is argued that star formation could provide most of the energy. Interestingly, most of these objects appear to be in interacting systems, suggesting a triggering mechanism. Exploration of this population to higher redshift will have particular significance for models of AGN/galaxy evolution.

3 The ELAIS Survey

The *ISO* observations took place between 12th March 1996 and 8th April 1998, shortly before the Helium boil off. The total survey area covered in blank fields is 6, 11, 12, & 1 sq. deg. at 6.75, 15, 90 and 175 μm , an additional 2 sq. deg. within the ELAIS survey areas has been surveyed by the FIRBACK team [4]. Table 1 specifies the main survey areas.

4 Comparison with other *ISO* surveys

ISO carried out a variety of surveys exploring the available parameter space of depth and area. Table 2 summaries the main extra-galactic blank-field surveys.

Table 2: Field Surveys with *ISO*

Survey Name	[e.g. ref]	Wavelength / μm	Integration /s	Area /sqdeg
PHT Serendipity Survey	[1]	175	0.5	7000
CAM Parallel Mode	[16]	7	150	33
ELAIS		7,15,90,175	40, 40, 24, 128	6, 11, 12,1
CAM Shallow	[6]	15	180	1.3
FIR Back	[4]	175	256, 128	1, 3
IR Back	[10]	90, 135,180	23, 27, 27	1, 1, 1
SA 57	[11]	60, 90	150, 50	0.42,0.42
CAM Deep	[6]	7, 15, 90	800, 990, 144	0.28, 0.28, 0.28
Comet fields	[3]	12	302	0.11
CFRS	[8]	7,15,60,90	720, 1000, 3000,3000	0.067,0.067,0.067,0.067
CAM Ultra-Deep	[6]	7	3520	0.013
ISOHDF South	[13]	7, 15	> 6400, > 6400	4.7e-3, 4.7e-3
Deep SSA13	[17]	7	34000	2.5e-3
Deep Lockman	[9]	7, 90, 175	44640, 48, 128	2.5e-3, 1.2 , 1
ISOHDF North	[14]	7, 15	12800, 6400	1.4e-3, 4.2e-3

5 Provisional Number Counts

Source catalogues have been extracted from the ELAIS data at all wavelengths and a very provisional source count analysis has been performed at 15 μm . The results of this are shown in Figure 1. As can be seen these counts confirm the strong evolution detected in the *ISO HDF* analysis [12]. A more detailed analysis of the counts will be presented shortly [15]

6 Follow-up

An extensive follow-up programme is being undertaken, including measurements at all wavelengths from X-ray to radio [2]. This programme will provide essential information for identifying the types of objects detected in the infrared, their luminosities, energy budgets and other detailed properties. As well as studying the properties of the objects detected by *ISO* a number of the follow-up surveys will provide independent source lists which will be extremely valuable in their own right, not least because we can investigate why some objects emit in the infrared and others do not.

Acknowledgements. This paper is based on observations with *ISO*, an ESA project, with instruments funded by ESA Member States (especially the PI coun-




Figure 1: Provisional 15 μm integral source counts from the ELAIS survey, together counts from *ISO HDF* ([12]) with *IRAS* 12 μm counts (thick line) at bright end, (Rush *et al.* 1993 *IRAS* data shifted to 15 μm using cirrus spectrum). Models are from [7]: all components solid; spiral galaxies dotted; star-bursts dash-dot-dot-dot; s0 dot-dash; AGN-dash Elliptical galaxies-short dashes. Faint end constraints (hatched) come from the *ISO HDF P(D)* analysis. The depths probed by some other *ISO* surveys are indicated CAM-D & CAM-S [5].

tries: France, Germany, the Netherlands and the United Kingdom) and with participation of ISAS and NASA. This work was in part supported by PPARC grant no. GR/K98728 and EC Network is FMRX-CT96-0068

References

- [1] Bogun, S., *et al.*, (1996), *A&A*, 315L, 71
- [2] Ciliegì P. *et al.* 1998 MNRAS (in press) astro-ph/9805353
- [3] Clements D., *et al.* (in prep.)
- [4] Clements D., *et al.* 1998 astro-ph/9809054
- [5] Elbaz D., 1997, ‘Taking *ISO* to the Limits’ Laureijs R. & Levine D., (ESA)
- [6] Elbaz D. *et al.*, 1998, astro-ph/9807209
- [7] Franceschini A., *et al.* 1994, ApJ, 427, 140
- [8] Hammer, F., Flores, H. astro-ph/9806184
- [9] Kawara, K. *et al.* 1998 A&A 336 9
- [10] Mattila *et al.* (in prep.)
- [11] Nørgaard-Nielsen H.U., *et al* 1997, ‘Taking *ISO* to the Limits’ Laureijs R. & Levine D. (ESA)
- [12] Oliver S.J. *et al*, 1997, MNRAS, 289, 471
- [13] Oliver S.J. *et al*, (in prep.)
- [14] Serjeant, S.B.G., *et al*, 1997, MNRAS, 289, 457
- [15] Serjeant, S.B.G., *et al*, (in prep.)
- [16] Siebenmorgen, R.; , *et al.*, 1996, *A&A*, 315L. 169
- [17] Taniguchi Y., *et al* 1997, ‘Taking *ISO* to the Limits’ Laureijs R. & Levine D., (ESA)